

## **General Disclaimer**

### **One or more of the Following Statements may affect this Document**

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

J 72-19301 e1  
CR-128548



# THE JOHNS HOPKINS UNIVERSITY

DEPARTMENT  
OF  
PHYSICS

## SPECIAL REPORT

NASA/MSC Contract NAS 9-11528  
Task I. Principal Investigator Services

A BRIEF HISTORY OF DATA AND  
THEORY PERTAINING TO THE  
ATMOSPHERE OF THE MOON

Submitted by:

G. E. Thomas  
University of Colorado  
Co-Investigator  
Apollo 17 UV Experiment

TECHNICAL LIBRARY  
BUILDING 45

NOV 8 1972

Manned Spacecraft Center  
Houston, Texas 77058

May 5, 1972

Baltimore, Maryland 21218

THE JOHNS HOPKINS UNIVERSITY

DEPARTMENT OF PHYSICS

HOMewood CAMPUS

BALTIMORE, MARYLAND 21218

May 5, 1972

To: Distribution

From: Wm. G. Fastie, Principal Investigator

Subject: Special Report on a Brief History of Data and Theory  
Pertaining to the Atmosphere of the Moon.

We submit herewith a special report prepared by  
G. E. Thomas, University of Colorado (co-investigator).  
This report presents some information about the atmosphere  
of the moon and includes a bibliography on the subject.

*Wm G Fastie*  
Wm. G. Fastie

WGF/eb

Encl.

Distribution:

5 Copies	NASA/MSC,	L. McFadin
1 Copy	" "	M. Krisberg
1 Copy	" "	E. Jones
1 Copy	" "	N. Hardee
1 Copy	" "	E. Smith
1 Copy	" "	H. Harvey
1 Copy	ONRRR/JHU	W. B. Girkin
1 Copy	JHU	A. P. Ashton
1 Copy	APL/JHU	T. Wyatt
1 Copy	APL/JHU NASARR	D. Beeman
1 Copy	NASA	G. F. Esenwein, Jr.
1 Copy	NASA/GSFC	J. Diggins
1 Copy	Univ. of Pitts	T. M. Donahue
1 Copy	Univ. of Colo.	G. E. Thomas
1 Copy	Univ. of Colo.	C. A. Barth
1 Copy	JHU	P. D. Feldman
1 Copy	JHU	R. C. Henry
1 Copy	NASA/MSC,	Code BT4

To remove, tear along this line

**A BRIEF HISTORY OF DATA AND  
THEORY PERTAINING TO THE ATMOSPHERE  
OF THE MOON**

**Submitted by:**

**G. E. Thomas**

**Co-Investigator**

**Apollo 17 UV Experiment**

**April 23, 1972**

## I. Introduction

This report consists of a brief historical resume of data and theory for the lunar atmosphere existing prior to the Apollo program. It is intended to be a summary of the most important pre-Apollo research, rather than an exhaustive bibliographical study, and constitutes a supporting analytical study for the Apollo 17 UV Spectrometer Experiment.

## II. Experimental Data on the Lunar Atmosphere

All Earth-based experiments designed to measure atmospheric density on the Moon have led only to upper limits. The first attempts by early astronomers to detect twilight effects or the bending of light during stellar occultations met with failure, and led only to estimates of upper limits of about  $10^{-4}$  atm (Opik, 1962). Efforts to detect lunar auroras (Khan, 1946) have always given negative results. This has led to an estimation of the upper limit on lunar density to be that of the Earth's atmosphere in the altitude range where auroras occur (100 - 120 km implying a density range of  $\sim 10^{-7}$  -  $10^{-8}$  atm, according to present knowledge). However this limit implicitly assumes the same energetic particle flux impinging on the Moon as in the Earth's auroral zones. The lack of a reliable detection of lunar meteors (although see the discussion by Opik (1962) of the observations of W. H. Haas) leads to a more definite upper limit corresponding to an altitude on Earth of

about 90 km (where the density is  $\sim 10^{-5}$  atm).

More sensitive tests for a lunar atmosphere were introduced by Fessenkopf (1943) who placed an upper limit of  $1.3 \times 10^{-7}$  atm, on the basis of a lack of polarization on the dark side of the lunar terminator. This conclusion was strongly criticized by Lyot and Dollfus (1940) because of Fessenkopf's assumption that the ashen light was unpolarized. Dollfus (1952) using a Savart-Lyot polariscope at Pic-du-Midi showed that the lack of polarization in the twilight zones near the poles implied an upper limit of  $10^{-9}$  atm. Using an improved technique he later extended this limit to  $10^{-10}$  atm (Dollfus, 1956). A lack of extension of the terminator at the edge of the visible disk led Opik (1955) to place an upper limit of  $10^{-6}$  atm. A widely-quoted observation by Kozyrev (1959) of a spectrogram of an "eruption" from the central peak of the crater Alphonsus points out the possibility of occasional local enhancements of atmosphere density by volcanism.

Gringauz et al (1961) reported direct measurements of ion densities at 10 Moon radii exceeding the interplanetary density, with an indication of an increasing density with decreasing altitude. Recent measurements using radio astronomy techniques have been reported by Elsmore (1957a, 1957b), Risbeth and Little (1957), Hazard et al (1963), and Pomalaza-Diaz (1967) which involve the refraction of signals from radio

star sources or from spacecraft during lunar occultations. The earlier occultation measurements indicated ion densities less than  $10^2 - 10^3 \text{ cm}^{-3}$ . The most accurate to date are from measurements of radio signals from Pioneer VII indicating less than  $40 \text{ ions-cm}^{-3}$  (Pomalaza-Diaz, 1967). A deduction of neutral density must be made from considerations of ionization sources and losses. The degree of ionization has sometimes been assumed to be that in the Earth's atmosphere near the F-2 peak ( $1 \times 10^{-3}$ ), which gave  $10^5 - 10^6 \text{ cm}^{-3}$  for the neutral gas (Ellsmore, 1957b). However as Johnson (1972) has pointed out, this degree of ionization is unrealistically high, as lunar ions have a much shorter lifetime than do those in the Earth's ionosphere. Depending upon the photo-ionization time and the loss times due to the ion being swept up by the magnetic field in the solar wind, the degree of ionization varies between  $5 \times 10^{-7}$  for atomic hydrogen (H) to  $2 \times 10^{-4}$  for Argon(A). Thus more modest upper limits must be placed on the lunar atmosphere from such experiments. For H, this upper limit is therefore about  $8 \times 10^7 \text{ cm}^{-3}$ , not much smaller than the upper limit of  $\sim 2 \times 10^9 \text{ cm}^{-3}$  imposed by Dollfus, (1956) optical limits.

### III. Theoretical Studies of the Lunar Atmosphere

Before the discovery of the solar wind, most speculations on the origin of a lunar atmosphere centered about the possibility of outgassing from the lunar interior, volatilization



of meteors, and radioactivity decay (for example the product of radioactive potassium  $K^{40}$  is  $A^{40}$ .) The primeval atmosphere was considered largely in the light of evaporative loss processes. Thus on the basis of the time constants for thermal escape from an exosphere (Spitzer, 1952, Biutner, 1958, 1959), it was generally expected that since for the rare gases Krypton (Kr) and Xenon (Xe) (mean atomic weights 83.8 and 131.3 respectively) the depletion time exceeded the lunar age of  $5 \times 10^9$  years, these gases would be the principal constituents of the lunar atmosphere. However such calculations refer to an average lunar surface temperature of  $400^\circ K$ . Opik and Singer (1960) pointed out that the expected pressure of  $10^{-13}$  atmosphere of Xe and Kr (Edwards and Borst, 1958) would lead to an atmosphere in which the temperature is comparable to the exospheric temperature of Earth ( $\sim 1500^\circ K$ ). This higher temperature would reduce the escape times to 170 years and  $4 \times 10^6$  years for Kr and Xe respectively. This result would indicate that the lunar atmosphere would adjust itself to a density ( $< 10^{14}$  atoms-cm<sup>-2</sup>-column) low enough for its temperature to be that of the surface itself; in other words the entire atmosphere is an exosphere. Opik and Singer also calculated the effect of the loss of photoelectrons produced at the surface by solar extreme ultraviolet. They estimated that an electric potential of 20-25 volts (positive) would exist at



the lunar surface, and that most  $\text{Kr}^+$  and  $\text{Xe}^+$  ions formed in the region where the potential is not yet screened would pick up sufficient energy (2.4 and 3.7 ev respectively) for escape. For an estimated screening thickness of 4 meters the lifetimes of Kr or Xe atoms were estimated to be about 3000 years. Thus it became clear that an independent lunar atmosphere (either of primeval origin or due to slow outgassing from the interior) cannot exist because of non-thermal loss processes even for the heaviest gases.

The discovery of the solar wind led to the possibility of an atmosphere formed from solar wind accretion (Gold, 1959; Opik, 1962; Nakada and Mihalov, 1962; Weil and Barasch, 1963; Bernstein et al, 1963; Hinton and Taeusch, 1964; Michel, 1964; Vogel, 1966) and an additional loss process from solar-wind "scavenging" (Herring and Licht, 1959; and above references). The usual method is to use the known (or parameterized) solar wind flux, assume cosmic abundance of the rare gases in the solar wind, and from a balance of all gains and losses, to extract the average surface density. The most complete study was that of Hinton and Taeusch who considered the following losses: photoionization and subsequent escape, thermal escape (which is the most important loss for hydrogen and helium), and charge exchange and impact ionization from the

solar wind. The sources assumed by Hinton and Taeusch were either solar wind or volcanism (the latter scaled from the known rates for Earth).

#### IV. Recent Theoretical and Experimental Results

Quantitative studies of the horizontal and vertical distribution of the lunar atmosphere began with Hodges and Johnson (1968), who showed that due to lateral transport, the horizontal variation of gases heavier than helium goes like  $T_s^{-5/2}$ , where  $T_s$  is the surface temperature. More recently Hodges (1972) has extended this work in terms of a generalized diffusion model. Gott and Potter (1968) presented detailed calculations of H and H<sub>2</sub> distributions in a hypothetical lunar atmosphere, and the Lyman alpha brightness expected from resonance scattering of solar Lyman alpha emission.

Very recently, there have appeared two papers, the first a review paper by Johnson (1971) who calculates surface densities from solar wind accretion using recent knowledge of solar wind fluxes (Bame et al, 1970) and ionization rates. Using the  $T_s^{-5/2}$  law of Hodges and Johnson (1968) he predicts that the daytime lunar atmosphere is composed of Ne ( $6 \times 10^4 \text{ cm}^{-3}$ ), He ( $6 \times 10^3 \text{ cm}^{-3}$ ), H ( $5 \times 10^3 \text{ cm}^{-3}$ ) and A ( $1.2 \times 10^3 \text{ cm}^{-3}$ ) with traces of Kr ( $8 \text{ cm}^3$ ) and Xe ( $1.3 \text{ cm}^{-3}$ ). The nighttime densities, because of the  $T_s^{-5/2}$  law for the heavier constituents,

are expected to be much greater for Ne ( $1.5 \times 10^6 \text{ cm}^{-3}$ ) and A ( $3 \times 10^4 \text{ cm}^{-3}$ ). H and He densities do not follow the above law since they do not satisfy the assumption that the average horizontal distance travelled in the exosphere is small. However the behaviour should still be inversely dependent on temperature. A very different boundary condition for the H and He fluxes was given by Vogel (1966), who assumed a steady daytime flux varying as the cosine of the solar zenith angle. Yeh and Chang (1972) have recently used this as a lower boundary condition and calculated the three-dimensional spatial variation of H in the lunar atmosphere. As expected this yields a surface density that decreases into the night-side, by a factor of  $\sim 4$ . However a realistic lateral transport model for H and He has yet to be calculated.

Hodges and Johnson (1968) pointed out that such gases as Kr and Xe, which liquefy at temperatures encountered on the nightside, are subject to surface adsorption causing them to essentially vanish from the nighttime lunar atmosphere. Rapid heating at sunrise would release most of these adsorbed gases producing a sharp maximum near the sunrise terminator. This expectation was not borne out by actual measurements through sunrise by the cold cathode gauge experiment placed on the Moon by Apollo 14 (Johnson et al, 1972). Instead of a gradual rise in density as the sun rose, there appeared a

very steep rise in density from the nighttime value of  $2 \times 10^5 \text{ cm}^{-3}$  to  $\sim 10^7 \text{ cm}^{-3}$ . The daytime value is believed to be a result of solar-induced outgassing of some contaminant in the vicinity of the landing area. The low nighttime density is attributed by Johnson et al to the fact that the surface material is not saturated with neon. (In other words, the Moon is a sink of neon atoms given off by the sun.) This possibility was suggested in a more general context by Gutkin et al (1969).

## References

- Bame, S. J., J. R. Ashbridge, A. J. Hundhausen, and M. D. Montgomery, "Solar Wind Ions," J. Geophys. Res., 75, 6360, 1970.
- Bernstein, W. R., W. Fredericks, J. L. Vogl, and W. A. Fowler, "The Lunar Atmosphere and the Solar Wind," Icarus, 2, 233, 1963.
- Biutner, E. K., "The Dissipation of Gas from Planetary Atmospheres. I. The Dissipation of Isothermal Ideal Gas in a Central Field of Gravitation," Astron. Zh., 35, 572, 1958.
- Biutner, E. K., "The Dissipation of Gas from Planetary Atmospheres. II. The Total Velocity of Dissipation of Gas from a Planetary Atmosphere. The Problem of Terrestrial Helium," Astron. Zh., 36, 89, 1959.
- Brandt, J. C., "Density of the Lunar Atmosphere," Science, 131, 1606, 1960.
- Dollfus, A., Compt. Rendu. Acad. Sci., 234, 2046, 1952.
- Dollfus, A., Ann. Astrophys., 19, 83, 1956.
- Edwards, W. F. and L. B. Borst, "Possible Sources of Lunar Atmosphere," Science, 127, 325, 1968.
- Ellsmore, B., "Radio Observation of the Lunar Atmosphere," Phil. Mag., 2, 1040, 1957b.

Elsmore, B., "The Lunar Occultation of a Radio Star and the Derivation of an Upper Limit for the Density of a Lunar Atmosphere," Radio Astronomy, 47-49, Ed. H. C. Vande Hulst, Cambridge Univ. Press, London, 1957a.

Fessenkopf, V. G., "On the Mass of the Moon's Atmosphere," Astr. J. Soviet Union, 20, 1, 1943.

Firsoff, V., "Dissipation of Planetary Atmospheres," Science, 130, 1337, 1959.

Gold, T., In discussion following R. Jastrow, "Outer Atmospheres of the Earth and Planets," J. Geophys. Res., 64, 1798, 1959.

Gott, R. and A. E. Potter, Jr., "Distribution of Atomic Hydrogen in the Lunar Atmosphere," Trans. Am. Geophys. Union, 49, 706, 1968.

Gringauz, K. I., V. V. Bezrukikh, V. D. Ozerov, and R. E. Rybchinskii, "A Study of Interplanetary Ionized Gases, Energetic Electrons, and Solar Corpuscular Emission," Artificial Earth Satellites, 122-129, Plenum Press, New York, 1961.

Gutkin, A. M., M. S. Markov, C. M. Raitburd, and M. V. Slonimskaya, "Adsorptive Mechanism for Loss of the Moon's Atmosphere," Defense Scientific Information Service Translation, T514R, May 1969 (Transl. from Dok. Akad. Nauk SSSR, 182, 1968).

- Hazard, C., M. B. Mackey, and A. J. Shimmins, "Investigation of the Radio Source 3C 273 by the Method of Lunar Occultations," Nature, 198, 1037, 1963.
- Herring, J. R. and A. L. Licht, "Effect of the Solar Wind on the Lunar Atmosphere," Science, 130, 266, 1959.
- Hinton, F. L. and D. R. Taeusch, "Variation of the Lunar Atmosphere with the Strength of the Solar Wind," J. Geophys. Res., 73, 7307, 1968.
- Hodges, R. R., Jr., "Applicability of a Diffusion Model to Lateral Transport in the Terrestrial and Lunar Exospheres," Planet. Space Sci., 20, 103, 1972.
- Hodges, R. R., Jr. and F. S. Johnson, "Lateral Transport in Planetary Exospheres," J. Geophys. Res., 73, 7307, 1968.
- Johnson, F. S., D. E. Evans, and J. M. Carroll, "Observations of Lunar Atmosphere," Space Res. XII, 1972.
- Johnson, F. S., "Lunar Atmosphere," Rev. Geophys. Space Phys., 9, 813, 1971.
- Khan, M. A. R., "A Decisive Test for the Presence of even a Highly Rarefied Lunar Atmosphere," Pop. Astr., 54, 312, 1946.
- Kozyrev, N. A., "Priroda, 84, March 1959; Sky and Telesc., 18, 561, 1959.
- Lyot, B. and A. Dollfus, "Recherche d'une Atmosphere au Voisinage de la lune," Compt. Rendu. Acad. Sci., 229, 1277, 1949.



- Michel, F. C., "Interaction between the Solar Wind and the Lunar Ionosphere," Planet. Space Sci., 12, 1075, 1964.
- Nakada, M. P. and J. D. Mihalov, "Accretion of Solar Wind from a Lunar Atmosphere," J. Geophys. Res., 67, 1670, 1962.
- Opik, E. J., Irish Astron. J., 3, 137, 1955.
- Opik, E. J. and S. F. Singer, "Escape of Gases from the Moon," J. Geophys. Res., 65, 3065, 1960.
- Opik, E. J., "The Lunar Atmosphere," Planet. Space Sci., 9, 211, 1962.
- Pomalaza-Diaz, J. C., "Measurement of the Lunar Ionosphere by Occultation of the Pioneer VII Spacecraft," Stanford Electronics Laboratories Scientific Report No. SU-SEL-67-095, 1967.
- Risbeth, H. and A. G. Little, Observatory, 77, 897, 1957.
- Singer, S. F., "Atmosphere Near the Moon," Astronaut. Acta, 7, 135, 1961.
- Singer, S. F., "Atmosphere Near the Moon," Conf. Lunar Exploration, Virginia Poly Tech. Inst., Blacksburg, Va. IV-1 to IV-11, 1962.
- Spitzer, L. J., The Terrestrial Atmosphere above 300 km in "The Atmospheres of the Earth and Planets," Ed. G. P. Kuiper, Univ. of Chicago Press, 1957.
- Vestine, E. H., "Evolution and Nature of the Lunar Atmosphere," RAND Corp. Res. Mem., RM-2106, January 1958.

Vogel, U., "Molecular Fluxes in the Lunar Atmosphere," Planet. Space Sci., 14, 1233, 1966.

Watson, K., B. C. Murray, and H. Brown, "The Behaviour of Volatiles on the Lunar Surface," J. Geophys. Res., 66, 3033, 1961.

Weil, H. and M. L. Barasch, "A Theoretical Lunar Ionosphere," Icarus, 1, 346, 1963.

Weil, N. A., "Lunar and Planetary Surface Conditions," Acad. Press, New York, 1965.

Yeh, T. T. J. and G. K. Chang, "Density and Flux Distributions of Neutral Gases in the Lunar Atmosphere," J. Geophys. Res., 77, 1720, 1972.